

< Back to results | 1 of 2 Next >

Export

Download

Print

E-mail

Save to PDF

Add to List



More... >

Full Text

View at Publisher

Iranian Journal of Medical Physics Open Access
Volume 15, Issues 1-3, October 2018, Pages 271-276

A phantom study for the optimization of image quality and radiation dose for common radiographic examinations in digital radiography (Article)

Moey, S.-F. , Shazli, Z.A. 
Department of Diagnostic Imaging and Radiotherapy, Kulliyah (Faculty) of Allied Health Sciences, International Islamic University Malaysia, Kuantan Campus, Kuantan, Pahang, 25200, Malaysia

Abstract View references (14)

Introduction: Phantom studies facilitate the implementation of radiation dose surveillance as a function of radiographic technical parameters for minimizing patient radiation dose. The evidence of such investigations can then be used to evaluate technical parameters used in the radiographic procedures to reduce radiation dose without compromising the image quality. Material and Methods: This experimental study was carried out using an anthropomorphic phantom and the Leeds test object. Computed radiographic system was utilized and the images were printed for objective evaluation. Dose-area-product (DAP) readings were obtained using a DAP meter for the technical parameters employed for the radiographic procedures. Results: The use of 0.2 mm additional copper filtration resulted in the lowest radiation doses for all four radiographic procedures (i.e. posteroanterior chest, anteroposterior abdomen and lumbar sacral spine projections). The highest tube potential appropriate to the body part being imaged, patient size, image receptor response and required information resulted in the minimum radiation dose to the patient without compromising the image quality. The focus to film distance utilized for the radiographic procedure must be in accordance with the focus to grid distance specified by the manufacturer when using the bucky to eliminate grid "cut-off." Conclusion: The optimization of image quality and radiation dose can be accomplished by using a phantom and selecting the imaging parameters that yield an acceptable image quality with the lowest entrance surface dose while considering the adjustment for patient size. © 2018 Mashhad University of Medical Sciences.

SciVal Topic Prominence ⓘ

Topic: Radiographic Image Enhancement | Radiography | radiography CR

Prominence percentile: 66.986 ⓘ

Author keywords

Digital radiography Radiation dosage Radiation protection Radiologic phantom

Indexed keywords


Engineering controlled terms: Dosimetry E-learning Image quality Phantoms Quality control Radiation Radiography

Engineering uncontrolled terms: Anthropomorphic phantoms Digital radiography Imaging parameters Objective evaluation Patient radiation dose Radiation dosage Radiographic examination Radiographic systems

Metrics ⓘ

0 Citations in Scopus
0 Field-Weighted Citation Impact



PlumX Metrics 
Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

Related documents

Measurement of X-ray for the dose area product and spectrum by the added filtration in Rando phantom

Lee, J.-W. , Kim, E.-S. , Choi, J. (2014) *Radiation Effects and Defects in Solids*

Increased filtration and image receptor exposure

Sandridge, T.G. (2016) *Radiologic Technology*

The effect of compensating filter on image quality in lateral projection of thoraco lumbar radiography

Daud, N.A.A. , Ali, M.H. , Nazri, N.A.A. (2014) *Journal of Physics: Conference Series*

View all related documents based on references

Find more related documents in Scopus based on:

NEW! SciVal Topic Prominence is now available in Scopus.

Which Topic is this article related to? View the Topic.



ISSN: 1735160X

Source Type: Journal

Original language: English



DOI: 10.22038/ijmp.2018.29825.1325

Document Type: Article

Publisher: Mashhad University of Medical Sciences

References (14)

[View in search results format >](#)

☐ All [Export](#)  Print  E-mail [Save to PDF](#) [Create bibliography](#)

- ☐ 1 (2004) *Managing patient dose in digital radiology*. Cited 20 times.
ICRP Publication 93. Ann. ICRP 34 (1)

- ☐ 2 (1996) *Radiological protection and safety in medicine*. Cited 123 times.
ICRP Publication 73. Ann. ICRP 26 (2)

- ☐ 3 Thali, M.J., Viner, M.D., Brogdon, B.G.
Brogdon's Forensic Radiology. Cited 52 times.
2nd ed. 2010. CRC Press: New York

- ☐ 4 Karami, V., Zabihzadeh, M., Keshtkar, M.
Evaluation of the Entrance Surface Dose (ESD) and radiation dose to the radiosensitive organs in pediatric pelvic radiography

(2017) *International Journal of Pediatrics*, 5 (6), pp. 5013-5022. Cited 2 times.
http://ijp.mums.ac.ir/article_8382_0d0ee3c6cba635a7639785c98f9b67a2.pdf
doi: 10.22038/ijp.2017.22364.1873

[View at Publisher](#)

- ☐ 5 Joseph, D., Igashi, J., Shen, S., Eshiet, P., Yabwa, D., Lutsi, G., Mundi, A., (...), Joseph, G.
Assessment of entrance skin dose and image quality of chest x-rays in two university teaching hospitals in north east Nigeria
(2014) *IOSR Journal of Nursing and Health Science*, 3 (6).
(II)

- ☐ 6 de Souza, A.B.B., Dias, S.K., Augusto, F.M., Marques, W.A.
Evaluation of entrance surface air kerma rate and clinical images quality in chest radiography

(2009) *Radiologia Brasileira*, 42 (5), pp. 315-319. Cited 2 times.
http://www.scielo.br/pdf/rb/v42n5/en_a11v42n5.pdf

[View at Publisher](#)